

Acute Lymphocytic Leukemia Interphase Chromosome Profiling (ALL-ICP): A Simple and Comprehensive Method to Detect Chromosome Abnormalities in ALL Samples

R. Babu¹, D.L. Van Dyke², J. Tepperberg³, E. Fuentes¹, S. Fuentes¹, S. Kopuri¹, C. Williamson², M. Liu⁴, S. Papa¹, S. Schwartz³, P. Papenhausen³, P. Koduru⁵

¹Department of Research and Development, InteGen LLC, Orlando, FL 32819; ²Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, MN 55902; ³Department of Cytogenetics, Lab Corp of America, Research Triangle Park, NC 27709; ⁴Department of Clinical Cytogenetics, Genetics Associates Inc., Nashville, TN 37203 ⁵Department of Pathology, UT Southwestern Medical Center, Dallas, TX 75390

INTRODUCTION

B- and T-cell acute lymphocytic leukemia (ALL) affects both children and adults. Cytogenetic and recent molecular genetic findings at diagnosis constitute important, independent prognostic factors in all age groups. Due to the typically poor morphology of the chromosome preparations, conventional cytogenetic analysis can miss one or more recurrent abnormalities and most ALL FISH panels only target select abnormalities. Therefore, a simple, fast, and comprehensive improvement to karyotype analysis would greatly enhance the diagnostic capabilities of clinical service as well as research laboratories for ALL.

MATERIALS & METHODS

We recently developed and validated a novel molecular technology, Interphase Chromosome Profiling (ICP) (Babu et al., 2017). We tested a variation (ALL-ICP) of our original method for interrogating interphase cells and metaphase chromosomes in ALL samples. The design consisted of a multiplex approach with analysis of six chromosomes per hybridization site. Telomeres and pericentromeric regions on each chromosome are targeted and the resulting fluorescent signals are spectrally distinct and easily recognizable from each other using two dual filter sets from Chroma. An additional probe set targeting common deletions and duplication/amplification was included. Overnight hybridization is done on one slide in four areas and whenever possible 20 metaphase spreads are analyzed for each chromosome. When 20 metaphases are unavailable, interphase nuclei were used to complete the 20-cell analysis. Modified culture conditions included increased time and exposure to dilute concentrations of Colcemid and/or exposure to a chemical known to induce premature chromosome condensation during the last three hours of culture. Forty-eight ALL samples were studied blindly by the laboratory that developed the technology. Three institutions provided cell pellets with known cytogenetic and FISH results.

RESULTS

Sample Number	Cytogenetic Findings	ICP Results	Additional changes found and/or characterized by ALL -ICP		
1	Hyperdiploid, +der	Concordant +	Characterized the derivative		
2	Add(5q), inv(7)	Inv missed	Add origin 15q		
	Tetraploid, add(3p), 6(p)				
3	1-3 mar	Concordant +	Deletion 14q, markers origin 13q and 10c		
4	Complex abnormalities	Concordant +	14q deletion and dim 14q		
5	Complex, add and	Missed interstitial	Add origin 19q		
	interstitial deletions	deletions	Add ongin 194		
	Complex, add(5q),	Concordant +,	Add origin 5q and 8q resulting in		
6	add(17q), interstitial	Missed interstitial	duplications		
	deletions	deletions	·		
7	Complex with markers	Concordant +	One mar is der(11)		
8	-X and 6q deletion – 1 cell	Concordant +	t(12;21) and 14q deletion		
9	Hyperdiploid, i(7q)	Concordant	None		
10	Hyperdiploid, add(19p)	Concordant +	Add origin 12q		
11	Hyperdiploid	Concordant +	+9 is der(9)t(8;9); +15 is +18		
12	dic(9;20),+21	Concordant	None		
13	t(1;19)(q23;p13.3)	Concordant +	14q deletion		
14	t(1;19)(q23;p13.3)	Concordant	None		
		Concordant +			
15	t(9;22)(q34;q11.2)		Homozygous 14q deletion		
16	t(9;22)(q34;q11.2)	Concordant +	14q deletion		
17	+i(21)(q10)x2	Concordant +	t(5;10), 16q and 22q deletions		
18	Hyperdiploid	Concordant +	+14 is der(14)t(14;21), deletions		
			(subtelomere) 18p and 15q		
<u> 19</u>	del(13)(q14q34)	Concordant +	t(12;21), 14q deletion		
20	Hypodiploid, t(9;22),	Missed duplication	None		
	dup(3q)	Wilssed daplication	None		
21	Hyperdiploid with dup(1q)	Missed duplication	del(3p) and 13 identified as der(3); deletion 16p and -20		
22	t(6;22)(q23;q12)	Concordant +	t(12;21) and multiple rearrangements involving chromosomes 6, 9, 8, 10		
23	Complex with variant t(9;22)	Concordant +	12p deletion		
24	add(21)(q22)	Concordant +	21q deletion, RUNX1 duplication		
25	+X,+21c	Concordant +	21q deletion, 14q deletion, der(15)		
26	del (12)(p11.2p13)	Concordant	None		
27	Variant t(9;22)	Concordant +	Iso dic(17q)		
28	del(6)(q13q27)	Concordant	None		
29	Complex with add(10q)	Concordant +	Clarified add(10q) as iso dic(10q)		
30	Deletions 9p and 6q	Concordant +	-19		
31	add (6q)	Concordant +	Add origin 5q; der(19)t(1;19), 14q		
			deletion		
32	Hyperdiploid	Concordant +	10p and 21q deletions		
33	9p (<i>CDKN2A</i>) deletion	Concordant	None		
34	12p deletion; add	Concordant+	t(12;21), add material from chromosome		
35	Complex; FISH positive for	Concordant+	X and 11 t(12;21), 14q32 homozygous deletion, 8p deletion, 6q deletion, <i>RUNX1</i>		
	ETV6-RUNX1		amplification		
36	t(7;17)	Concordant+	CDKN2A dim (partial deletion), t(1;7)		
37	t(2;14), +10	Concordant+	11q deletion; homozygous 14q deletion		
38	t(9;22)	Concordant+	Homozygous <i>CDKN2A</i> deletion; 12p deletion, 14q deletion, 4p deletion		
39	Complex karyotype; 'add' chromosomes	Concordant+	Add material from chromosomes 13q, 3p 11/9q		
40	Hyperdiploid; t(9;22); cyto	Concordant+	ICP 20 cells abnormal		
41	Hyperdiploid plus mars;	Concordant+	mars are copies of extra chromosomes; TCF3 deletion (19p13.3) missed by ICP –		
42	FISH 19p deletion 6q deletion in two cells;	Concordant+	not targeted 6q deletion in 20 cells		
	9p homozygous deletion 45,XY,dic(20;21)(q13.3;q2		dic t(20;21) has tandem duplication of		
43	2)	Concordant+	RUNX1 as well as telomere; 20q deletion confirmed		
44	9q deletion in 80% of the cells Marker chromosome -12:	Concordant+	9p deletion; balanced t(5;11)		
45	Marker chromosome, -12; ETV6-RUNX1 by FISH	Concordant+	Marker is possible variant translocation t(12;19;21)		
46	46,XX,t(2;7)(p21;p22)	Concordant+	14q deletion; possible t(2;12)		
47	Hyperdiploid	Concordant+	Possible 12p deletion		
48	Hyperdiploid	Concordant	None		

Concordant

None

IMAGES





FIGURE LEGEND

A – C: Representative images showing a 3-fold increase in mitotic index


- D: Trisomy 4 and 10
- E: Normal chromosomes 2, 14 and 22 Mix 1
- F: Normal chromosomes 8, 9 and 11 Mix 1
- G: 14q32 (*IGH*) deletion
- H: Homozygous 9p deletion and 6q deletion
- I: der(6)(5;6)(q;q)
- J: RUNX1 Amplification
- K L: t(12;21)(p13;q22)

DESIGN









CHROMOSOME COLOR SCHEME

		Mix 3	Mix 4	Mix 5
2	1	4	7	9p21.3
14	12	10	16	6q21
22	17	Υ	13	13q14.2
8	X	5	3	21q22.1 2
9	19	6	18	Xp22.33
11	21	20	15	
	14 22 8 9	14 12 22 17 8 X 9 19	14 12 10 22 17 Y 8 X 5 9 19 6	14 12 10 16 22 17 Y 13 8 X 5 3 9 19 6 18

CONCLUSION

- All numerical abnormalities identified by karyotype or FISH are detected by ALL-ICP.
- ALL-ICP refined the breakpoints and clarified marker and derivative chromosomes.
- Novel 14q32 (IgH) deletions and several previously unknown sub-telomere deletions were identified.
- There appears to be an association between t(9;22) and 14q deletions nearly 50% with this deletion had the translocation.
- ALL common deletions and amplification of RUNX1 were detected.
- Non-recurring interstitial deletions and duplications not targeted by ALL-ICP were not detected.
- All known and new balanced translocations were detected.
- ALL-ICP detected additional abnormalities in 85% of the cases.
- A three-fold increase in mitotic index was observed in modified cultures.

ALL-ICP is a simple method that saves valuable time and detects all clinically relevant recurrent B- and T-cell chromosome abnormalities. Additionally, it characterizes the marker chromosomes and 'add' material. The improved diagnostic capabilities, supported by the study results, make ALL-ICP ideal for the genetic diagnostic workup of Acute Lymphocytic Leukemia on standard as well as modified cytogenetic harvests.

REFERENCES & ACKNOWLEDGEMENTS

- 1) Interphase Chromosome Profiling: A Method for Conventional Banded Chromosome Analysis Using Interphase Nuclei. Babu R, Van Dyke DL, Dev VG et al., Arch Pathol Lab Med. doi: 10.5858/ arpa.2016-0621-OA
- 2) Clinical Significance of Cytogenetic Abnormalities in Adult Acute Lymphoblastic Leukemia. Faderl S, Kantarjian HM, Talpaz M, and Estrov Z. Blood, Vol 91, No 11 (June 1), 1998: pp 3995-4019
- 3) Cytogenetics and Molecular Genetics of Acute Lymphoblastic Leukemia. . Mrózek1 K, Harper DP, and Aplan PD. Hematol Oncol Clin North Am. 2009 October ; 23(5): 991–v. doi:10.1016/j.hoc.2009.07.001
- 4) Microdeletions are a general feature of adult and adolescent acute lymphoblastic leukemia:
- MacDougall F et al., PNAS , vol. 105, no. 18 , May 6, 2008: 6708–6713

5) Unexpected similarities with pediatric disease. Paulsson K, Cazier JB, Finlay

We would like to thank Yvonne Banol (InteGen LLC) for technical support.

Please contact us at contact@integenllc.com www.integenllc.com



48 Hyperdiploid